

AMENDMENTS TO THE CLAIMS:

1-17. (cancelled)

18. (currently amended) A method of treating morning stiffness, loss of grip strength, painful joints, or swollen joints in a patient suffering from morning stiffness, loss of grip strength, painful joints, or swollen joints, consisting of comprising identifying the patient suffering from morning stiffness, loss of grip strength, painful joints, or swollen joints and administering to the patient suffering from morning stiffness, loss of grip strength, painful joints, or swollen joints a morning stiffness, loss of grip strength, painful joints, or swollen joints treating effective amount of erythropoietin over a treatment period.

19. (canceled)

20. (currently amended) A method of ameliorating an erythrocyte sedimentation rate or C-reactive protein level in a patient in need of such amelioration, consisting of comprising identifying the patient in need of such amelioration and administering to the patient an erythrocyte sedimentation rate or C-reactive protein level activity ameliorating effective amount of erythropoietin over a period.

21-22. (canceled)

23. (previously presented) The method of claim 18, wherein the erythropoietin is human erythropoietin.

24. (previously presented) The method of claim 18, wherein the erythropoietin is of recombinant origin.

25. (previously presented) The method of claim 20, wherein the erythropoietin is human erythropoietin.

26. (previously presented) The method of claim 20, wherein the erythropoietin is of recombinant origin.

27-30. (canceled)

31. (previously presented) The method of claim 20, wherein the period comprises 6 weeks of treatment.

32. (previously presented) The method of claim 18 wherein the patient suffers from rheumatoid arthritis.

33. (previously presented) The method of claim 20 wherein the patient suffers from rheumatoid arthritis.

34. (currently amended) The method of claim 18 wherein the treatment period is at least 3 to 6 weeks.

35. (currently amended) The method of claim 20 wherein the treatment period is at least 3 to 6 weeks.